



Molecular Cell Biology

EIGHTH EDITION

Harvey Lodish

Arnold Berk

Chris A. Kaiser

Monty Krieger

Anthony Bretscher

Hidde Ploegh

Angelika Amon

Kelsey C. Martin



w.h.freeman

Macmillan Learning

New York

CONTENTS IN BRIEF

Part I Chemical and Molecular Foundations

- 1 Molecules, Cells, and Model Organisms 1
- 2 Chemical Foundations 31
- 3 Protein Structure and Function 67
- 4 Culturing and Visualizing Cells 129

Part II Biomembranes, Genes, and Gene Regulation

- 5 Fundamental Molecular Genetic Mechanisms 167
- 6 Molecular Genetic Techniques 223
- 7 Biomembrane Structure 271
- 8 Genes, Genomics, and Chromosomes 301
- 9 Transcriptional Control of Gene Expression 353
- 10 Post-transcriptional Gene Control 417

Part III Cellular Organization and Function

- 11 Transmembrane Transport of Ions and Small Molecules 473
- 12 Cellular Energetics 513
- 13 Moving Proteins into Membranes and Organelles 583
- 14 Vesicular Traffic, Secretion, and Endocytosis 631
- 15 Signal Transduction and G Protein–Coupled Receptors 673
- 16 Signaling Pathways That Control Gene Expression 719
- 17 Cell Organization and Movement I: Microfilaments 775
- 18 Cell Organization and Movement II: Microtubules and Intermediate Filaments 821
- 19 The Eukaryotic Cell Cycle 873

Part IV Cell Growth and Differentiation

- 20 Integrating Cells into Tissues 921
- 21 Stem Cells, Cell Asymmetry, and Cell Death 975
- 22 Cells of the Nervous System 1025
- 23 Immunology 1079
- 24 Cancer 1135

CONTENTS

Preface

vii

Part I Chemical and Molecular Foundations

1 Molecules, Cells, and Model Organisms 1

1.1 The Molecules of Life 5

Proteins Give Cells Structure and Perform Most Cellular Tasks 7

Nucleic Acids Carry Coded Information for Making Proteins at the Right Time and Place 7

Phospholipids Are the Conserved Building Blocks of All Cellular Membranes 9

1.2 Prokaryotic Cell Structure and Function 10

Prokaryotes Comprise Two Kingdoms: Archaea and Eubacteria 10

Escherichia coli Is Widely Used in Biological Research 11

1.3 Eukaryotic Cell Structure and Function 12

The Cytoskeleton Has Many Important Functions 12

The Nucleus Contains the DNA Genome, RNA Synthetic Apparatus, and a Fibrous Matrix 12

Eukaryotic Cells Contain a Large Number of Internal Membrane Structures 14

Mitochondria Are the Principal Sites of ATP Production in Aerobic Cells 18

Chloroplasts Contain Internal Compartments in Which Photosynthesis Takes Place 18

All Eukaryotic Cells Use a Similar Cycle to Regulate Their Division 18

1.4 Unicellular Eukaryotic Model Organisms 19

Yeasts Are Used to Study Fundamental Aspects of Eukaryotic Cell Structure and Function 19

Mutations in Yeast Led to the Identification of Key Cell Cycle Proteins 21

Studies in the Alga *Chlamydomonas reinhardtii* Led to the Development of a Powerful Technique to Study Brain Function 22

The Parasite That Causes Malaria Has Novel Organelles That Allow It to Undergo a Remarkable Life Cycle 22

1.5 Metazoan Structure, Differentiation, and Model Organisms 24

Multicellularity Requires Cell-Cell and Cell-Matrix Adhesions 24

Epithelia Originated Early in Evolution 24

Tissues Are Organized into Organs 24

Genomics Has Revealed Important Aspects of Metazoan Evolution and Cell Function 24

Embryonic Development Uses a Conserved Set of Master Transcription Factors 25

Planaria Are Used to Study Stem Cells and Tissue Regeneration 27

Invertebrates, Fish, Mice, and Other Organisms Serve as Experimental Systems for Study of Human Development and Disease 28

Genetic Diseases Elucidate Important Aspects of Cell Function 28

The Following Chapters Present Much Experimental Data That Explains How We Know What We Know About Cell Structure and Function 29

2 Chemical Foundations 31

2.1 Covalent Bonds and Noncovalent Interactions 33

The Electronic Structure of an Atom Determines the Number and Geometry of the Covalent Bonds It Can Make 33

Electrons May Be Shared Equally or Unequally in Covalent Bonds 34

Covalent Bonds Are Much Stronger and More Stable Than Noncovalent Interactions 36

Ionic Interactions Are Attractions Between Oppositely Charged Ions 36

Hydrogen Bonds Are Noncovalent Interactions That Determine the Water Solubility of Uncharged Molecules 37

Van der Waals Interactions Are Weak Attractive Interactions Caused by Transient Dipoles 38

The Hydrophobic Effect Causes Nonpolar Molecules to Adhere to One Another	39	NAD ⁺ and FAD Couple Many Biological Oxidation and Reduction Reactions	63
Molecular Complementarity Due to Noncovalent Interactions Leads to a Lock-and-Key Fit Between Biomolecules	40		
2.2 Chemical Building Blocks of Cells	41	3 Protein Structure and Function	67
Amino Acids Differing Only in Their Side Chains Compose Proteins	42	3.1 Hierarchical Structure of Proteins	69
Five Different Nucleotides Are Used to Build Nucleic Acids	45	The Primary Structure of a Protein Is Its Linear Arrangement of Amino Acids	69
Monosaccharides Covalently Assemble into Linear and Branched Polysaccharides	46	Secondary Structures Are the Core Elements of Protein Architecture	70
Phospholipids Associate Noncovalently to Form the Basic Bilayer Structure of Biomembranes	48	Tertiary Structure Is the Overall Folding of a Polypeptide Chain	72
		There Are Four Broad Structural Categories of Proteins	72
2.3 Chemical Reactions and Chemical Equilibrium	51	Different Ways of Depicting the Conformation of Proteins Convey Different Types of Information	74
A Chemical Reaction Is in Equilibrium When the Rates of the Forward and Reverse Reactions Are Equal	52	Structural Motifs Are Regular Combinations of Secondary Structures	75
The Equilibrium Constant Reflects the Extent of a Chemical Reaction	52	Domains Are Modules of Tertiary Structure	76
Chemical Reactions in Cells Are at Steady State	52	Multiple Polypeptides Assemble into Quaternary Structures and Supramolecular Complexes	78
Dissociation Constants of Binding Reactions Reflect the Affinity of Interacting Molecules	53	Comparing Protein Sequences and Structures Provides Insight into Protein Function and Evolution	79
Biological Fluids Have Characteristic pH Values	54		
Hydrogen Ions Are Released by Acids and Taken Up by Bases	55	3.2 Protein Folding	81
Buffers Maintain the pH of Intracellular and Extracellular Fluids	55	Planar Peptide Bonds Limit the Shapes into Which Proteins Can Fold	81
		The Amino Acid Sequence of a Protein Determines How It Will Fold	81
2.4 Biochemical Energetics	57	Folding of Proteins in Vivo Is Promoted by Chaperones	82
Several Forms of Energy Are Important in Biological Systems	57	Protein Folding Is Promoted by Proline Isomerases	86
Cells Can Transform One Type of Energy into Another	58	Abnormally Folded Proteins Can Form Amyloids That Are Implicated in Diseases	87
The Change in Free Energy Determines If a Chemical Reaction Will Occur Spontaneously	58		
The ΔG° of a Reaction Can Be Calculated from Its K_{eq}	60	3.3 Protein Binding and Enzyme Catalysis	89
The Rate of a Reaction Depends on the Activation Energy Necessary to Energize the Reactants into a Transition State	60	Specific Binding of Ligands Underlies the Functions of Most Proteins	89
Life Depends on the Coupling of Unfavorable Chemical Reactions with Energetically Favorable Ones	61	Enzymes Are Highly Efficient and Specific Catalysts	90
Hydrolysis of ATP Releases Substantial Free Energy and Drives Many Cellular Processes	61	An Enzyme's Active Site Binds Substrates and Carries Out Catalysis	91
ATP Is Generated During Photosynthesis and Respiration	62	Serine Proteases Demonstrate How an Enzyme's Active Site Works	92
		Enzymes in a Common Pathway Are Often Physically Associated with One Another	96
		3.4 Regulating Protein Function	97
		Regulated Synthesis and Degradation of Proteins Is a Fundamental Property of Cells	97
		The Proteasome Is a Molecular Machine Used to Degrade Proteins	97

Ubiquitin Marks Cytosolic Proteins for Degradation in Proteasomes	99	Growth of Cells in Two-Dimensional and Three-Dimensional Culture Mimics the In Vivo Environment	133
Noncovalent Binding Permits Allosteric, or Cooperative, Regulation of Proteins	100	Hybridomas Produce Abundant Monoclonal Antibodies	135
Noncovalent Binding of Calcium and GTP Are Widely Used as Allosteric Switches to Control Protein Activity	101	A Wide Variety of Cell Biological Processes Can Be Studied with Cultured Cells	136
Phosphorylation and Dephosphorylation Covalently Regulate Protein Activity	102	Drugs Are Commonly Used in Cell Biological Research	136
Ubiquitinylation and Deubiquitinylation Covalently Regulate Protein Activity	103		
Proteolytic Cleavage Irreversibly Activates or Inactivates Some Proteins	104		
Higher-Order Regulation Includes Control of Protein Location	105		
3.5 Purifying, Detecting, and Characterizing Proteins	105	4.2 Light Microscopy: Exploring Cell Structure and Visualizing Proteins Within Cells	139
Centrifugation Can Separate Particles and Molecules That Differ in Mass or Density	106	The Resolution of the Conventional Light Microscope Is About 0.2 μm	139
Electrophoresis Separates Molecules on the Basis of Their Charge-to-Mass Ratio	107	Phase-Contrast and Differential-Interference-Contrast Microscopy Visualize Unstained Live Cells	141
Liquid Chromatography Resolves Proteins by Mass, Charge, or Affinity	109	Imaging Subcellular Details Often Requires That Specimens Be Fixed, Sectioned, and Stained	142
Highly Specific Enzyme and Antibody Assays Can Detect Individual Proteins	111	Fluorescence Microscopy Can Localize and Quantify Specific Molecules in Live Cells	143
Radioisotopes Are Indispensable Tools for Detecting Biological Molecules	114	Intracellular Ion Concentrations Can Be Determined with Ion-Sensitive Fluorescent Dyes	143
Mass Spectrometry Can Determine the Mass and Sequence of Proteins	116	Immunofluorescence Microscopy Can Detect Specific Proteins in Fixed Cells	144
Protein Primary Structure Can Be Determined by Chemical Methods and from Gene Sequences	118	Tagging with Fluorescent Proteins Allows the Visualization of Specific Proteins in Live Cells	146
Protein Conformation Is Determined by Sophisticated Physical Methods	119	Deconvolution and Confocal Microscopy Enhance Visualization of Three-Dimensional Fluorescent Objects	147
		Two-Photon Excitation Microscopy Allows Imaging Deep into Tissue Samples	149
		TIRF Microscopy Provides Exceptional Imaging in One Focal Plane	150
3.6 Proteomics	122	FRAP Reveals the Dynamics of Cellular Components	151
Proteomics Is the Study of All or a Large Subset of Proteins in a Biological System	122	FRET Measures Distance Between Fluorochromes	152
Advanced Techniques in Mass Spectrometry Are Critical to Proteomic Analysis	123	Super-Resolution Microscopy Can Localize Proteins to Nanometer Accuracy	153
		Light-Sheet Microscopy Can Rapidly Image Cells in Living Tissue	155
4 Culturing and Visualizing Cells	129		
4.1 Growing and Studying Cells in Culture	130	4.3 Electron Microscopy: High-Resolution Imaging	156
Culture of Animal Cells Requires Nutrient-Rich Media and Special Solid Surfaces	130	Single Molecules or Structures Can Be Imaged Using a Negative Stain or Metal Shadowing	157
Primary Cell Cultures and Cell Strains Have a Finite Life Span	131	Cells and Tissues Are Cut into Thin Sections for Viewing by Electron Microscopy	158
Transformed Cells Can Grow Indefinitely in Culture	132	Immunoelectron Microscopy Localizes Proteins at the Ultrastructural Level	159
Flow Cytometry Separates Different Cell Types	132	Cryoelectron Microscopy Allows Visualization of Specimens Without Fixation or Staining	160

Scanning Electron Microscopy of Metal-Coated Specimens Reveals Surface Features 161

4.4 Isolation of Cell Organelles 161

Disruption of Cells Releases Their Organelles and Other Contents 162

Centrifugation Can Separate Many Types of Organelles 162

Organelle-Specific Antibodies Are Useful in Preparing Highly Purified Organelles 162

Proteomics Reveals the Protein Composition of Organelles 164

Part II Biomembranes, Genes, and Gene Regulation

5 Fundamental Molecular Genetic Mechanisms 167

5.1 Structure of Nucleic Acids 169

A Nucleic Acid Strand Is a Linear Polymer with End-to-End Directionality 170

Native DNA Is a Double Helix of Complementary Antiparallel Strands 170

DNA Can Undergo Reversible Strand Separation 172

Torsional Stress in DNA Is Relieved by Enzymes 174

Different Types of RNA Exhibit Various Conformations Related to Their Functions 174

5.2 Transcription of Protein-Coding Genes and Formation of Functional mRNA 176

A Template DNA Strand Is Transcribed into a Complementary RNA Chain by RNA Polymerase 176

Organization of Genes Differs in Prokaryotic and Eukaryotic DNA 179

Eukaryotic Precursor mRNAs Are Processed to Form Functional mRNAs 180

Alternative RNA Splicing Increases the Number of Proteins Expressed from a Single Eukaryotic Gene 181

5.3 The Decoding of mRNA by tRNAs 183

Messenger RNA Carries Information from DNA in a Three-Letter Genetic Code 183

The Folded Structure of tRNA Promotes Its Decoding Functions 185

Nonstandard Base Pairing Often Occurs Between Codons and Anticodons 186

Amino Acids Become Activated When Covalently Linked to tRNAs 188

5.4 Stepwise Synthesis of Proteins on Ribosomes 188

Ribosomes Are Protein-Synthesizing Machines 188

Methionyl-tRNA^{Met} Recognizes the AUG Start Codon 190

Eukaryotic Translation Initiation Usually Occurs at the First AUG Downstream from the 5' End of an mRNA 191

During Chain Elongation Each Incoming Aminoacyl-tRNA Moves Through Three Ribosomal Sites 193

Translation Is Terminated by Release Factors When a Stop Codon Is Reached 195

Polysomes and Rapid Ribosome Recycling Increase the Efficiency of Translation 195

GTPase-Superfamily Proteins Function in Several Quality-Control Steps of Translation 195

Nonsense Mutations Cause Premature Termination of Protein Synthesis 196

5.5 DNA Replication 197

DNA Polymerases Require a Primer to Initiate Replication 197

Duplex DNA Is Unwound, and Daughter Strands Are Formed at the DNA Replication Fork 199

Several Proteins Participate in DNA Replication 199

DNA Replication Occurs Bidirectionally from Each Origin 201

5.6 DNA Repair and Recombination 203

DNA Polymerases Introduce Copying Errors and Also Correct Them 203

Chemical and Radiation Damage to DNA Can Lead to Mutations 203

High-Fidelity DNA Excision-Repair Systems Recognize and Repair Damage 204

Base Excision Repairs T-G Mismatches and Damaged Bases 205

Mismatch Excision Repairs Other Mismatches and Small Insertions and Deletions 205

Nucleotide Excision Repairs Chemical Adducts that Distort Normal DNA Shape 206

Two Systems Use Recombination to Repair Double-Strand Breaks in DNA 207

Homologous Recombination Can Repair DNA Damage and Generate Genetic Diversity 209

5.7 Viruses: Parasites of the Cellular Genetic System 212

Most Viral Host Ranges Are Narrow 212

Viral Capsids Are Regular Arrays of One or a Few Types of Protein 213

Viruses Can Be Cloned and Counted in Plaque Assays	213
Lytic Viral Growth Cycles Lead to Death of Host Cells	213
Viral DNA Is Integrated into the Host-Cell Genome in Some Nonlytic Viral Growth Cycles	216

6 Molecular Genetic Techniques 223

6.1 Genetic Analysis of Mutations to Identify and Study Genes 224

Recessive and Dominant Mutant Alleles Generally Have Opposite Effects on Gene Function	224
Segregation of Mutations in Breeding Experiments Reveals Their Dominance or Recessivity	225
Conditional Mutations Can Be Used to Study Essential Genes in Yeast	227
Recessive Lethal Mutations in Diploids Can Be Identified by Inbreeding and Maintained in Heterozygotes	228
Complementation Tests Determine Whether Different Recessive Mutations Are in the Same Gene	229
Double Mutants Are Useful in Assessing the Order in Which Proteins Function	230
Genetic Suppression and Synthetic Lethality Can Reveal Interacting or Redundant Proteins	231
Genes Can Be Identified by Their Map Position on the Chromosome	232

6.2 DNA Cloning and Characterization 234

Restriction Enzymes and DNA Ligases Allow Insertion of DNA Fragments into Cloning Vectors	234
Isolated DNA Fragments Can Be Cloned into <i>E. coli</i> Plasmid Vectors	236
Yeast Genomic Libraries Can Be Constructed with Shuttle Vectors and Screened by Functional Complementation	237
cDNA Libraries Represent the Sequences of Protein-Coding Genes	238
The Polymerase Chain Reaction Amplifies a Specific DNA Sequence from a Complex Mixture	239
Cloned DNA Molecules Can Be Sequenced Rapidly by Methods Based on PCR	243

6.3 Using Cloned DNA Fragments to Study Gene Expression 246

Hybridization Techniques Permit Detection of Specific DNA Fragments and mRNAs	246
DNA Microarrays Can Be Used to Evaluate the Expression of Many Genes at One Time	247
Cluster Analysis of Multiple Expression Experiments Identifies Co-regulated Genes	248
<i>E. coli</i> Expression Systems Can Produce Large Quantities of Proteins from Cloned Genes	249

Plasmid Expression Vectors Can Be Designed for Use in Animal Cells	251
--	-----

6.4 Locating and Identifying Human Disease Genes 254

Monogenic Diseases Show One of Three Patterns of Inheritance	254
DNA Polymorphisms Are Used as Markers for Linkage Mapping of Human Mutations	255
Linkage Studies Can Map Disease Genes with a Resolution of About 1 Centimorgan	256
Further Analysis Is Needed to Locate a Disease Gene in Cloned DNA	257
Many Inherited Diseases Result from Multiple Genetic Defects	257

6.5 Inactivating the Function of Specific Genes in Eukaryotes 259

Normal Yeast Genes Can Be Replaced with Mutant Alleles by Homologous Recombination	260
Genes Can Be Placed Under the Control of an Experimentally Regulated Promoter	260
Specific Genes Can Be Permanently Inactivated in the Germ Line of Mice	261
Somatic Cell Recombination Can Inactivate Genes in Specific Tissues	261
Dominant-Negative Alleles Can Inhibit the Function of Some Genes	262
RNA Interference Causes Gene Inactivation by Destroying the Corresponding mRNA	264
Engineered CRISPR–Cas9 Systems Allow Precise Genome Editing	266

7 Biomembrane Structure 271

7.1 The Lipid Bilayer: Composition and Structural Organization 273

Phospholipids Spontaneously Form Bilayers	273
Phospholipid Bilayers Form a Sealed Compartment Surrounding an Internal Aqueous Space	274
Biomembranes Contain Three Principal Classes of Lipids	276
Most Lipids and Many Proteins Are Laterally Mobile in Biomembranes	278
Lipid Composition Influences the Physical Properties of Membranes	279
Lipid Composition Is Different in the Exoplasmic and Cytosolic Leaflets	281
Cholesterol and Sphingolipids Cluster with Specific Proteins in Membrane Microdomains	282
Cells Store Excess Lipids in Lipid Droplets	283

7.2 Membrane Proteins: Structure and Basic Functions	284		
Proteins Interact with Membranes in Three Different Ways	284		
Most Transmembrane Proteins Have Membrane-Spanning α Helices	285		
Multiple β Strands in Porins Form Membrane-Spanning "Barrels"	288		
Covalently Attached Lipids Anchor Some Proteins to Membranes	288		
All Transmembrane Proteins and Glycolipids Are Asymmetrically Oriented in the Bilayer	289		
Lipid-Binding Motifs Help Target Peripheral Proteins to the Membrane	290		
Proteins Can Be Removed from Membranes by Detergents or High-Salt Solutions	290		
7.3 Phospholipids, Sphingolipids, and Cholesterol: Synthesis and Intracellular Movement	293		
Fatty Acids Are Assembled from Two-Carbon Building Blocks by Several Important Enzymes	293		
Small Cytosolic Proteins Facilitate Movement of Fatty Acids	293		
Fatty Acids Are Incorporated into Phospholipids Primarily on the ER Membrane	294		
Flippases Move Phospholipids from One Membrane Leaflet to the Opposite Leaflet	295		
Cholesterol Is Synthesized by Enzymes in the Cytosol and ER Membrane	295		
Cholesterol and Phospholipids Are Transported Between Organelles by Several Mechanisms	296		
8 Genes, Genomics, and Chromosomes	301		
8.1 Eukaryotic Gene Structure	303		
Most Eukaryotic Genes Contain Introns and Produce mRNAs Encoding Single Proteins	303		
Simple and Complex Transcription Units Are Found in Eukaryotic Genomes	303		
Protein-Coding Genes May Be Solitary or Belong to a Gene Family	305		
Heavily Used Gene Products Are Encoded by Multiple Copies of Genes	307		
Nonprotein-Coding Genes Encode Functional RNAs	308		
8.2 Chromosomal Organization of Genes and Noncoding DNA	309		
Genomes of Many Organisms Contain Nonfunctional DNA	309		
Most Simple-Sequence DNAs Are Concentrated in Specific Chromosomal Locations	310		
		DNA Fingerprinting Depends on Differences in Length of Simple-Sequence DNAs	311
		Unclassified Intergenic DNA Occupies a Significant Portion of the Genome	312
		8.3 Transposable (Mobile) DNA Elements	312
		Movement of Mobile Elements Involves a DNA or an RNA Intermediate	313
		DNA Transposons Are Present in Prokaryotes and Eukaryotes	314
		LTR Retrotransposons Behave Like Intracellular Retroviruses	316
		Non-LTR Retrotransposons Transpose by a Distinct Mechanism	318
		Other Retroposed RNAs Are Found in Genomic DNA	321
		Mobile DNA Elements Have Significantly Influenced Evolution	321
		8.4 Genomics: Genome-Wide Analysis of Gene Structure and Function	323
		Stored Sequences Suggest Functions of Newly Identified Genes and Proteins	324
		Comparison of Related Sequences from Different Species Can Give Clues to Evolutionary Relationships Among Proteins	325
		Genes Can Be Identified Within Genomic DNA Sequences	326
		The Number of Protein-Coding Genes in an Organism's Genome Is Not Directly Related to Its Biological Complexity	326
		8.5 Structural Organization of Eukaryotic Chromosomes	327
		Chromatin Exists in Extended and Condensed Forms	328
		Modifications of Histone Tails Control Chromatin Condensation and Function	330
		Nonhistone Proteins Organize Long Chromatin Loops	335
		Additional Nonhistone Proteins Regulate Transcription and Replication	339
		8.6 Morphology and Functional Elements of Eukaryotic Chromosomes	341
		Chromosome Number, Size, and Shape at Metaphase Are Species-Specific	341
		During Metaphase, Chromosomes Can Be Distinguished by Banding Patterns and Chromosome Painting	341
		Chromosome Painting and DNA Sequencing Reveal the Evolution of Chromosomes	342

Interphase Polytene Chromosomes Arise by DNA Amplification	343
Three Functional Elements Are Required for Replication and Stable Inheritance of Chromosomes	345
Centromere Sequences Vary Greatly in Length and Complexity	345
Addition of Telomeric Sequences by Telomerase Prevents Shortening of Chromosomes	347

9 Transcriptional Control of Gene Expression 353

9.1 Control of Gene Expression in Bacteria 356

Transcription Initiation by Bacterial RNA Polymerase Requires Association with a Sigma Factor	357
Initiation of <i>lac</i> Operon Transcription Can Be Repressed or Activated	357
Small Molecules Regulate Expression of Many Bacterial Genes via DNA-Binding Repressors and Activators	358
Transcription Initiation from Some Promoters Requires Alternative Sigma Factors	359
Transcription by σ^{54} -RNA Polymerase Is Controlled by Activators That Bind Far from the Promoter	359
Many Bacterial Responses Are Controlled by Two-Component Regulatory Systems	360
Expression of Many Bacterial Operons Is Controlled by Regulation of Transcriptional Elongation	361

9.2 Overview of Eukaryotic Gene Control 363

Regulatory Elements in Eukaryotic DNA Are Found Both Close to and Many Kilobases Away from Transcription Start Sites	364
Three Eukaryotic RNA Polymerases Catalyze Formation of Different RNAs	367
The Largest Subunit in RNA Polymerase II Has an Essential Carboxy-Terminal Repeat	370

9.3 RNA Polymerase II Promoters and General Transcription Factors 371

RNA Polymerase II Initiates Transcription at DNA Sequences Corresponding to the 5' Cap of mRNAs	371
The TATA Box, Initiators, and CpG Islands Function as Promoters in Eukaryotic DNA	371
General Transcription Factors Position RNA Polymerase II at Start Sites and Assist in Initiation	373
Elongation Factors Regulate the Initial Stages of Transcription in the Promoter-Proximal Region	377

9.4 Regulatory Sequences in Protein-Coding Genes and the Proteins Through Which They Function 378

Promoter-Proximal Elements Help Regulate Eukaryotic Genes	378
Distant Enhancers Often Stimulate Transcription by RNA Polymerase II	379
Most Eukaryotic Genes Are Regulated by Multiple Transcription-Control Elements	379
DNase I Footprinting and EMSA Detect Protein-DNA Interactions	380
Activators Are Composed of Distinct Functional Domains	381
Repressors Are the Functional Converse of Activators	383
DNA-Binding Domains Can Be Classified into Numerous Structural Types	384
Structurally Diverse Activation and Repression Domains Regulate Transcription	386
Transcription Factor Interactions Increase Gene-Control Options	387
Multiprotein Complexes Form on Enhancers	388

9.5 Molecular Mechanisms of Transcription Repression and Activation 390

Formation of Heterochromatin Silences Gene Expression at Telomeres, near Centromeres, and in Other Regions	390
Repressors Can Direct Histone Deacetylation at Specific Genes	393
Activators Can Direct Histone Acetylation at Specific Genes	394
Chromatin-Remodeling Complexes Help Activate or Repress Transcription	395
Pioneer Transcription Factors Initiate the Process of Gene Activation During Cellular Differentiation	395
The Mediator Complex Forms a Molecular Bridge Between Activation Domains and Pol II	396

9.6 Regulation of Transcription-Factor Activity 398

DNase I Hypersensitive Sites Reflect the Developmental History of Cellular Differentiation	398
Nuclear Receptors Are Regulated by Extracellular Signals	400
All Nuclear Receptors Share a Common Domain Structure	400
Nuclear-Receptor Response Elements Contain Inverted or Direct Repeats	400
Hormone Binding to a Nuclear Receptor Regulates Its Activity as a Transcription Factor	402
Metazoans Regulate the RNA Polymerase II Transition from Initiation to Elongation	402
Termination of Transcription Is Also Regulated	402

9.7 Epigenetic Regulation of Transcription	404		
DNA Methylation Represses Transcription	404		
Methylation of Specific Histone Lysines Is Linked to Epigenetic Mechanisms of Gene Repression	405		
Epigenetic Control by Polycomb and Trithorax Complexes	406		
Long Noncoding RNAs Direct Epigenetic Repression in Metazoans	409		
9.8 Other Eukaryotic Transcription Systems	412		
Transcription Initiation by Pol I and Pol III Is Analogous to That by Pol II	412		
10 Post-transcriptional Gene Control	417		
10.1 Processing of Eukaryotic Pre-mRNA	419		
The 5' Cap Is Added to Nascent RNAs Shortly After Transcription Initiation	420		
A Diverse Set of Proteins with Conserved RNA-Binding Domains Associate with Pre-mRNAs	421		
Splicing Occurs at Short, Conserved Sequences in Pre-mRNAs via Two Transesterification Reactions	423		
During Splicing, snRNAs Base-Pair with Pre-mRNA	424		
Spliceosomes, Assembled from snRNPs and a Pre-mRNA, Carry Out Splicing	426		
Chain Elongation by RNA Polymerase II Is Coupled to the Presence of RNA-Processing Factors	428		
SR Proteins Contribute to Exon Definition in Long Pre-mRNAs	428		
Self-Splicing Group II Introns Provide Clues to the Evolution of snRNAs	429		
3' Cleavage and Polyadenylation of Pre-mRNAs Are Tightly Coupled	430		
Nuclear Exoribonucleases Degrade RNA That Is Processed Out of Pre-mRNAs	432		
RNA Processing Solves the Problem of Pervasive Transcription of the Genome in Metazoans	432		
10.2 Regulation of Pre-mRNA Processing	435		
Alternative Splicing Generates Transcripts with Different Combinations of Exons	435		
A Cascade of Regulated RNA Splicing Controls <i>Drosophila</i> Sexual Differentiation	435		
Splicing Repressors and Activators Control Splicing at Alternative Sites	437		
		RNA Editing Alters the Sequences of Some Pre-mRNAs	439
		10.3 Transport of mRNA Across the Nuclear Envelope	440
		Phosphorylation and Dephosphorylation of SR Proteins Imposes Directionality on mRNP Export Across the Nuclear Pore Complex	441
		Balbani Rings in Insect Larval Salivary Glands Allow Direct Visualization of mRNP Export Through NPCs	442
		Pre-mRNAs in Spliceosomes Are Not Exported from the Nucleus	443
		HIV Rev Protein Regulates the Transport of Unspliced Viral mRNAs	444
		10.4 Cytoplasmic Mechanisms of Post-transcriptional Control	445
		Degradation of mRNAs in the Cytoplasm Occurs by Several Mechanisms	445
		Adenines in mRNAs and lncRNAs May Be Post-transcriptionally Modified by N ⁶ Methylation	447
		Micro-RNAs Repress Translation and Induce Degradation of Specific mRNAs	447
		Alternative Polyadenylation Increases miRNA Control Options	450
		RNA Interference Induces Degradation of Precisely Complementary mRNAs	450
		Cytoplasmic Polyadenylation Promotes Translation of Some mRNAs	451
		Protein Synthesis Can Be Globally Regulated	452
		Sequence-Specific RNA-Binding Proteins Control Translation of Specific mRNAs	455
		Surveillance Mechanisms Prevent Translation of Improperly Processed mRNAs	456
		Localization of mRNAs Permits Production of Proteins at Specific Regions Within the Cytoplasm	457
		10.5 Processing of rRNA and tRNA	461
		Pre-rRNA Genes Function as Nucleolar Organizers	461
		Small Nucleolar RNAs Assist in Processing Pre-rRNAs	462
		Self-Splicing Group I Introns Were the First Examples of Catalytic RNA	466
		Pre-tRNAs Undergo Extensive Modification in the Nucleus	466
		Nuclear Bodies Are Functionally Specialized Nuclear Domains	468

Part III Cellular Organization and Function

11 Transmembrane Transport of Ions and Small Molecules 473

11.1 Overview of Transmembrane Transport 474

- Only Gases and Small Uncharged Molecules Cross Membranes by Simple Diffusion 474
- Three Main Classes of Membrane Proteins Transport Molecules and Ions Across Cellular Membranes 475

11.2 Facilitated Transport of Glucose and Water 477

- Uniport Transport Is Faster and More Specific than Simple Diffusion 477
- The Low K_m of the GLUT1 Uniporter Enables It to Transport Glucose into Most Mammalian Cells 478
- The Human Genome Encodes a Family of Sugar-Transporting GLUT Proteins 480
- Transport Proteins Can Be Studied Using Artificial Membranes and Recombinant Cells 480
- Osmotic Pressure Causes Water to Move Across Membranes 481
- Aquaporins Increase the Water Permeability of Cellular Membranes 481

11.3 ATP-Powered Pumps and the Intracellular Ionic Environment 483

- There Are Four Main Classes of ATP-Powered Pumps 484
- ATP-Powered Ion Pumps Generate and Maintain Ionic Gradients Across Cellular Membranes 485
- Muscle Relaxation Depends on Ca^{2+} ATPases That Pump Ca^{2+} from the Cytosol into the Sarcoplasmic Reticulum 486
- The Mechanism of Action of the Ca^{2+} Pump Is Known in Detail 486
- Calmodulin Regulates the Plasma-Membrane Pumps That Control Cytosolic Ca^{2+} Concentrations 489
- The Na^+/K^+ ATPase Maintains the Intracellular Na^+ and K^+ Concentrations in Animal Cells 489
- V-Class H^+ ATPases Maintain the Acidity of Lysosomes and Vacuoles 489
- ABC Proteins Export a Wide Variety of Drugs and Toxins from the Cell 491
- Certain ABC Proteins “Flip” Phospholipids and Other Lipid-Soluble Substrates from One Membrane Leaflet to the Other 493
- The ABC Cystic Fibrosis Transmembrane Regulator Is a Chloride Channel, Not a Pump 494

11.4 Nongated Ion Channels and the Resting Membrane Potential 495

- Selective Movement of Ions Creates a Transmembrane Electric Gradient 495
- The Resting Membrane Potential in Animal Cells Depends Largely on the Outward Flow of K^+ Ions Through Open K^+ Channels 497
- Ion Channels Are Selective for Certain Ions by Virtue of a Molecular “Selectivity Filter” 497
- Patch Clamps Permit Measurement of Ion Movements Through Single Channels 500
- Novel Ion Channels Can Be Characterized by a Combination of Oocyte Expression and Patch Clamping 501

11.5 Cotransport by Symporters and Antiporters 502

- Na^+ Entry into Mammalian Cells Is Thermodynamically Favored 502
- Na^+ -Linked Symporters Enable Animal Cells to Import Glucose and Amino Acids Against High Concentration Gradients 503
- A Bacterial $\text{Na}^+/\text{Amino Acid}$ Symporter Reveals How Symport Works 504
- A Na^+ -Linked Ca^{2+} Antiporter Regulates the Strength of Cardiac Muscle Contraction 504
- Several Cotransporters Regulate Cytosolic pH 505
- An Anion Antiporter Is Essential for Transport of CO_2 by Erythrocytes 506
- Numerous Transport Proteins Enable Plant Vacuoles to Accumulate Metabolites and Ions 507

11.6 Transcellular Transport 508

- Multiple Transport Proteins Are Needed to Move Glucose and Amino Acids Across Epithelia 508
- Simple Rehydration Therapy Depends on the Osmotic Gradient Created by Absorption of Glucose and Na^+ 509
- Parietal Cells Acidify the Stomach Contents While Maintaining a Neutral Cytosolic pH 509
- Bone Resorption Requires the Coordinated Function of a V-Class Proton Pump and a Specific Chloride Channel 510

12 Cellular Energetics 513

12.1 First Step of Harvesting Energy from Glucose: Glycolysis 515

- During Glycolysis (Stage I), Cytosolic Enzymes Convert Glucose to Pyruvate 516
- The Rate of Glycolysis Is Adjusted to Meet the Cell’s Need for ATP 516
- Glucose Is Fermented When Oxygen Is Scarce 518

12.2 The Structure and Functions of Mitochondria	520		
Mitochondria Are Multifunctional Organelles	520		
Mitochondria Have Two Structurally and Functionally Distinct Membranes	520		
Mitochondria Contain DNA Located in the Matrix	523		
The Size, Structure, and Coding Capacity of mtDNA Vary Considerably Among Organisms	525		
Products of Mitochondrial Genes Are Not Exported	526		
Mitochondria Evolved from a Single Endosymbiotic Event Involving a <i>Rickettsia</i> -Like Bacterium	527		
Mitochondrial Genetic Codes Differ from the Standard Nuclear Code	527		
Mutations in Mitochondrial DNA Cause Several Genetic Diseases in Humans	528		
Mitochondria Are Dynamic Organelles That Interact Directly with One Another	528		
Mitochondria Are Influenced by Direct Contacts with the Endoplasmic Reticulum	529		
12.3 The Citric Acid Cycle and Fatty Acid Oxidation	533		
In the First Part of Stage II, Pyruvate Is Converted to Acetyl CoA and High-Energy Electrons	533		
In the Second Part of Stage II, the Citric Acid Cycle Oxidizes the Acetyl Group in Acetyl CoA to CO ₂ and Generates High-Energy Electrons	533		
Transporters in the Inner Mitochondrial Membrane Help Maintain Appropriate Cytosolic and Matrix Concentrations of NAD ⁺ and NADH	535		
Mitochondrial Oxidation of Fatty Acids Generates ATP	536		
Peroxisomal Oxidation of Fatty Acids Generates No ATP	537		
12.4 The Electron-Transport Chain and Generation of the Proton-Motive Force	539		
Oxidation of NADH and FADH ₂ Releases a Significant Amount of Energy	539		
Electron Transport in Mitochondria Is Coupled to Proton Pumping	539		
Electrons Flow “Downhill” Through a Series of Electron Carriers	540		
Four Large Multiprotein Complexes Couple Electron Transport to Proton Pumping Across the Inner Mitochondrial Membrane	542		
The Reduction Potentials of Electron Carriers in the Electron-Transport Chain Favor Electron Flow from NADH to O ₂	546		
The Multiprotein Complexes of the Electron-Transport Chain Assemble into Supercomplexes	546		
Reactive Oxygen Species Are By-Products of Electron Transport	547		
		Experiments Using Purified Electron-Transport Chain Complexes Established the Stoichiometry of Proton Pumping	549
		The Proton-Motive Force in Mitochondria Is Due Largely to a Voltage Gradient Across the Inner Membrane	550
		12.5 Harnessing the Proton-Motive Force to Synthesize ATP	551
		The Mechanism of ATP Synthesis Is Shared Among Bacteria, Mitochondria, and Chloroplasts	552
		ATP Synthase Comprises F ₀ and F ₁ Multiprotein Complexes	553
		Rotation of the F ₁ γ Subunit, Driven by Proton Movement Through F ₀ , Powers ATP Synthesis	554
		Multiple Protons Must Pass Through ATP Synthase to Synthesize One ATP	555
		F ₀ c Ring Rotation Is Driven by Protons Flowing Through Transmembrane Channels	556
		ATP-ADP Exchange Across the Inner Mitochondrial Membrane Is Powered by the Proton-Motive Force	556
		The Rate of Mitochondrial Oxidation Normally Depends on ADP Levels	558
		Mitochondria in Brown Fat Use the Proton-Motive Force to Generate Heat	558
		12.6 Photosynthesis and Light-Absorbing Pigments	560
		Thylakoid Membranes in Chloroplasts Are the Sites of Photosynthesis in Plants	560
		Chloroplasts Contain Large DNAs Often Encoding More Than a Hundred Proteins	560
		Three of the Four Stages in Photosynthesis Occur Only During Illumination	561
		Photosystems Comprise a Reaction Center and Associated Light-Harvesting Complexes	563
		Photoelectron Transport from Energized Reaction-Center Chlorophyll <i>a</i> Produces a Charge Separation	564
		Internal Antennas and Light-Harvesting Complexes Increase the Efficiency of Photosynthesis	566
		12.7 Molecular Analysis of Photosystems	567
		The Single Photosystem of Purple Bacteria Generates a Proton-Motive Force but No O ₂	567
		Chloroplasts Contain Two Functionally and Spatially Distinct Photosystems	567
		Linear Electron Flow Through Both Plant Photosystems Generates a Proton-Motive Force, O ₂ , and NADPH	568
		An Oxygen-Evolving Complex Is Located on the Luminal Surface of the PSII Reaction Center	569
		Multiple Mechanisms Protect Cells Against Damage from Reactive Oxygen Species During Photoelectron Transport	570

Cyclic Electron Flow Through PSI Generates a Proton-Motive Force but No NADPH or O ₂	570
Relative Activities of Photosystems I and II Are Regulated	571

12.8 CO₂ Metabolism During Photosynthesis 573

Rubisco Fixes CO ₂ in the Chloroplast Stroma	573
Synthesis of Sucrose Using Fixed CO ₂ Is Completed in the Cytosol	573
Light and Rubisco Activase Stimulate CO ₂ Fixation	574
Photorespiration Competes with Carbon Fixation and Is Reduced in C ₄ Plants	576

13 Moving Proteins into Membranes and Organelles 583

13.1 Targeting Proteins To and Across the ER Membrane 585

Pulse-Chase Experiments with Purified ER Membranes Demonstrated That Secreted Proteins Cross the ER Membrane	586
A Hydrophobic N-Terminal Signal Sequence Targets Nascent Secretory Proteins to the ER	586
Cotranslational Translocation Is Initiated by Two GTP-Hydrolyzing Proteins	588
Passage of Growing Polypeptides Through the Translocon Is Driven by Translation	589
ATP Hydrolysis Powers Post-translational Translocation of Some Secretory Proteins in Yeast	591

13.2 Insertion of Membrane Proteins into the ER 593

Several Topological Classes of Integral Membrane Proteins Are Synthesized on the ER	593
Internal Stop-Transfer Anchor and Signal-Anchor Sequences Determine Topology of Single-Pass Proteins	594
Multipass Proteins Have Multiple Internal Topogenic Sequences	597
A Phospholipid Anchor Tethers Some Cell-Surface Proteins to the Membrane	598
The Topology of a Membrane Protein Can Often Be Deduced from Its Sequence	599

13.3 Protein Modifications, Folding, and Quality Control in the ER 601

A Preformed N-Linked Oligosaccharide Is Added to Many Proteins in the Rough ER	601
Oligosaccharide Side Chains May Promote Folding and Stability of Glycoproteins	602

Disulfide Bonds Are Formed and Rearranged by Proteins in the ER Lumen	603
Chaperones and Other ER Proteins Facilitate Folding and Assembly of Proteins	604
Improperly Folded Proteins in the ER Induce Expression of Protein-Folding Catalysts	606
Unassembled or Misfolded Proteins in the ER Are Often Transported to the Cytosol for Degradation	607

13.4 Targeting of Proteins to Mitochondria and Chloroplasts 608

Amphipathic N-Terminal Targeting Sequences Direct Proteins to the Mitochondrial Matrix	609
Mitochondrial Protein Import Requires Outer-Membrane Receptors and Translocons in Both Membranes	610
Studies with Chimeric Proteins Demonstrate Important Features of Mitochondrial Protein Import	612
Three Energy Inputs Are Needed to Import Proteins into Mitochondria	613
Multiple Signals and Pathways Target Proteins to Submitochondrial Compartments	613
Import of Chloroplast Stromal Proteins Is Similar to Import of Mitochondrial Matrix Proteins	617
Proteins Are Targeted to Thylakoids by Mechanisms Related to Bacterial Protein Translocation	617

13.5 Targeting of Peroxisomal Proteins 619

A Cytosolic Receptor Targets Proteins with an SKL Sequence at the C-Terminus to the Peroxisomal Matrix	619
Peroxisomal Membrane and Matrix Proteins Are Incorporated by Different Pathways	621

13.6 Transport Into and Out of the Nucleus 622

Large and Small Molecules Enter and Leave the Nucleus via Nuclear Pore Complexes	622
Nuclear Transport Receptors Escort Proteins Containing Nuclear-Localization Signals into the Nucleus	624
A Second Type of Nuclear Transport Receptor Escorts Proteins Containing Nuclear-Export Signals Out of the Nucleus	625
Most mRNAs Are Exported from the Nucleus by a Ran-Independent Mechanism	627

14 Vesicular Traffic, Secretion, and Endocytosis 631

14.1 Techniques for Studying the Secretory Pathway 634

Transport of a Protein Through the Secretory Pathway Can Be Assayed in Live Cells	634
---	-----

Yeast Mutants Define Major Stages and Many Components in Vesicular Transport	635	The Acidic pH of Late Endosomes Causes Most Receptor-Ligand Complexes to Dissociate	662
Cell-Free Transport Assays Allow Dissection of Individual Steps in Vesicular Transport	637	The Endocytic Pathway Delivers Iron to Cells Without Dissociation of the Transferrin–Transferrin Receptor Complex in Endosomes	663
14.2 Molecular Mechanisms of Vesicle Budding and Fusion	638	14.6 Directing Membrane Proteins and Cytosolic Materials to the Lysosome	665
Assembly of a Protein Coat Drives Vesicle Formation and Selection of Cargo Molecules	638	Multivesicular Endosomes Segregate Membrane Proteins Destined for the Lysosomal Membrane from Proteins Destined for Lysosomal Degradation	665
A Conserved Set of GTPase Switch Proteins Controls the Assembly of Different Vesicle Coats	639	Retroviruses Bud from the Plasma Membrane by a Process Similar to Formation of Multivesicular Endosomes	666
Targeting Sequences on Cargo Proteins Make Specific Molecular Contacts with Coat Proteins	641	The Autophagic Pathway Delivers Cytosolic Proteins or Entire Organelles to Lysosomes	667
Rab GTPases Control Docking of Vesicles on Target Membranes	641		
Paired Sets of SNARE Proteins Mediate Fusion of Vesicles with Target Membranes	642		
Dissociation of SNARE Complexes After Membrane Fusion Is Driven by ATP Hydrolysis	644		
14.3 Early Stages of the Secretory Pathway	645	15 Signal Transduction and G Protein–Coupled Receptors	673
COPII Vesicles Mediate Transport from the ER to the Golgi	645	15.1 Signal Transduction: From Extracellular Signal to Cellular Response	675
COPI Vesicles Mediate Retrograde Transport Within the Golgi and from the Golgi to the ER	647	Signaling Molecules Can Act Locally or at a Distance	675
Anterograde Transport Through the Golgi Occurs by Cisternal Maturation	648	Receptors Bind Only a Single Type of Hormone or a Group of Closely Related Hormones	676
		Protein Kinases and Phosphatases Are Employed in Many Signaling Pathways	676
		GTP-Binding Proteins Are Frequently Used in Signal Transduction Pathways as On/Off Switches	677
		Intracellular “Second Messengers” Transmit Signals from Many Receptors	678
		Signal Transduction Pathways Can Amplify the Effects of Extracellular Signals	679
		15.2 Studying Cell-Surface Receptors and Signal Transduction Proteins	681
		The Dissociation Constant Is a Measure of the Affinity of a Receptor for Its Ligand	681
		Binding Assays Are Used to Detect Receptors and Determine Their Affinity and Specificity for Ligands	681
		Near-Maximal Cellular Response to a Signaling Molecule Usually Does Not Require Activation of All Receptors	682
		Sensitivity of a Cell to External Signals Is Determined by the Number of Cell-Surface Receptors and Their Affinity for Ligand	683
		Hormone Analogs Are Widely Used as Drugs	683
		Receptors Can Be Purified by Affinity Chromatography Techniques	683
		Immunoprecipitation Assays and Affinity Techniques Can Be Used to Study the Activity of Signal Transduction Proteins	684
14.4 Later Stages of the Secretory Pathway	650		
Vesicles Coated with Clathrin and Adapter Proteins Mediate Transport from the <i>trans</i> -Golgi	651		
Dynamin Is Required for Pinching Off of Clathrin-Coated Vesicles	652		
Mannose 6-Phosphate Residues Target Soluble Proteins to Lysosomes	653		
Study of Lysosomal Storage Diseases Revealed Key Components of the Lysosomal Sorting Pathway	655		
Protein Aggregation in the <i>trans</i> -Golgi May Function in Sorting Proteins to Regulated Secretory Vesicles	655		
Some Proteins Undergo Proteolytic Processing After Leaving the <i>trans</i> -Golgi	656		
Several Pathways Sort Membrane Proteins to the Apical or Basolateral Region of Polarized Cells	657		
14.5 Receptor-Mediated Endocytosis	659		
Cells Take Up Lipids from the Blood in the Form of Large, Well-Defined Lipoprotein Complexes	659		
Receptors for Macromolecular Ligands Contain Sorting Signals That Target Them for Endocytosis	660		

15.3 G Protein–Coupled Receptors: Structure and Mechanism	686	Activated Phospholipase C Generates Two Key Second Messengers Derived from the Membrane Lipid Phosphatidylinositol 4,5-Bisphosphate	709
All G Protein–Coupled Receptors Share the Same Basic Structure	686	The Ca ²⁺ -Calmodulin Complex Mediates Many Cellular Responses to External Signals	713
Ligand-Activated G Protein–Coupled Receptors Catalyze Exchange of GTP for GDP on the α Subunit of a Heterotrimeric G Protein	689	DAG Activates Protein Kinase C	714
Different G Proteins Are Activated by Different GPCRs and In Turn Regulate Different Effector Proteins	691	Integration of Ca ²⁺ and cAMP Second Messengers Regulates Glycogenolysis	714
15.4 G Protein–Coupled Receptors That Regulate Ion Channels	693	Signal-Induced Relaxation of Vascular Smooth Muscle Is Mediated by a Ca ²⁺ -Nitric Oxide-cGMP-Activated Protein Kinase G Pathway	714
Acetylcholine Receptors in the Heart Muscle Activate a G Protein That Opens K ⁺ Channels	693	16 Signaling Pathways That Control Gene Expression 719	
Light Activates Rhodopsin in Rod Cells of the Eye	694	16.1 Receptor Serine Kinases That Activate Smads	722
Activation of Rhodopsin by Light Leads to Closing of cGMP-Gated Cation Channels	695	TGF- β Proteins Are Stored in an Inactive Form in the Extracellular Matrix	722
Signal Amplification Makes the Rhodopsin Signal Transduction Pathway Exquisitely Sensitive	696	Three Separate TGF- β Receptor Proteins Participate in Binding TGF- β and Activating Signal Transduction	722
Rapid Termination of the Rhodopsin Signal Transduction Pathway Is Essential for the Temporal Resolution of Vision	697	Activated TGF- β Receptors Phosphorylate Smad Transcription Factors	724
Rod Cells Adapt to Varying Levels of Ambient Light by Intracellular Trafficking of Arrestin and Transducin	698	The Smad3/Smad4 Complex Activates Expression of Different Genes in Different Cell Types	724
15.5 G Protein–Coupled Receptors That Activate or Inhibit Adenylyl Cyclase	699	Negative Feedback Loops Regulate TGF- β /Smad Signaling	725
Adenylyl Cyclase Is Stimulated and Inhibited by Different Receptor-Ligand Complexes	699	16.2 Cytokine Receptors and the JAK/STAT Signaling Pathway	726
Structural Studies Established How G _{αs} -GTP Binds to and Activates Adenylyl Cyclase	701	Cytokines Influence the Development of Many Cell Types	727
cAMP Activates Protein Kinase A by Releasing Inhibitory Subunits	701	Binding of a Cytokine to Its Receptor Activates One or More Tightly Bound JAK Protein Tyrosine Kinases	728
Glycogen Metabolism Is Regulated by Hormone-Induced Activation of PKA	702	Phosphotyrosine Residues Are Binding Surfaces for Multiple Proteins with Conserved Domains	730
cAMP-Mediated Activation of PKA Produces Diverse Responses in Different Cell Types	703	SH2 Domains in Action: JAK Kinases Activate STAT Transcription Factors	731
Signal Amplification Occurs in the cAMP-PKA Pathway	704	Multiple Mechanisms Down-Regulate Signaling from Cytokine Receptors	731
CREB Links cAMP and PKA to Activation of Gene Transcription	704	16.3 Receptor Tyrosine Kinases	734
Anchoring Proteins Localize Effects of cAMP to Specific Regions of the Cell	705	Binding of Ligand Promotes Dimerization of an RTK and Leads to Activation of Its Intrinsic Tyrosine Kinase	734
Multiple Mechanisms Suppress Signaling from the GPCR/cAMP/PKA Pathway	706	Homo- and Hetero-oligomers of Epidermal Growth Factor Receptors Bind Members of the Epidermal Growth Factor Family	735
15.6 G Protein–Coupled Receptors That Trigger Elevations in Cytosolic and Mitochondrial Calcium	708	Activation of the EGF Receptor Results in the Formation of an Asymmetric Active Kinase Dimer	736
Calcium Concentrations in the Mitochondrial Matrix, ER, and Cytosol Can Be Measured with Targeted Fluorescent Proteins	709	Multiple Mechanisms Down-Regulate Signaling from RTKs	737

16.4 The Ras/MAP Kinase Pathway	739	On Binding Delta, the Notch Receptor Is Cleaved, Releasing a Component Transcription Factor	761
Ras, a GTPase Switch Protein, Operates Downstream of Most RTKs and Cytokine Receptors	739	Matrix Metalloproteases Catalyze Cleavage of Many Signaling Proteins from the Cell Surface	763
Genetic Studies in <i>Drosophila</i> Identified Key Signal-Transducing Proteins in the Ras/MAP Kinase Pathway	739	Inappropriate Cleavage of Amyloid Precursor Protein Can Lead to Alzheimer's Disease	763
Receptor Tyrosine Kinases Are Linked to Ras by Adapter Proteins	741	Regulated Intramembrane Proteolysis of SREBPs Releases a Transcription Factor That Acts to Maintain Phospholipid and Cholesterol Levels	763
Binding of Sos to Inactive Ras Causes a Conformational Change That Triggers an Exchange of GTP for GDP	742		
Signals Pass from Activated Ras to a Cascade of Protein Kinases Ending with MAP Kinase	742	16.8 Integration of Cellular Responses to Multiple Signaling Pathways: Insulin Action	766
Phosphorylation of MAP Kinase Results in a Conformational Change That Enhances Its Catalytic Activity and Promotes Its Dimerization	744	Insulin and Glucagon Work Together to Maintain a Stable Blood Glucose Level	766
MAP Kinase Regulates the Activity of Many Transcription Factors Controlling Early Response Genes	745	A Rise in Blood Glucose Triggers Insulin Secretion from the β Islet Cells	767
G Protein–Coupled Receptors Transmit Signals to MAP Kinase in Yeast Mating Pathways	746	In Fat and Muscle Cells, Insulin Triggers Fusion of Intracellular Vesicles Containing the GLUT4 Glucose Transporter to the Plasma Membrane	767
Scaffold Proteins Separate Multiple MAP Kinase Pathways in Eukaryotic Cells	746	Insulin Inhibits Glucose Synthesis and Enhances Storage of Glucose as Glycogen	769
		Multiple Signal Transduction Pathways Interact to Regulate Adipocyte Differentiation Through PPAR γ , the Master Transcriptional Regulator	770
16.5 Phosphoinositide Signaling Pathways	748	Inflammatory Hormones Cause Derangement of Adipose Cell Function in Obesity	770
Phospholipase C γ Is Activated by Some RTKs and Cytokine Receptors	749		
Recruitment of PI-3 Kinase to Activated Receptors Leads to Synthesis of Three Phosphorylated Phosphatidylinositols	749		
Accumulation of PI 3-Phosphates in the Plasma Membrane Leads to Activation of Several Kinases	750		
Activated Protein Kinase B Induces Many Cellular Responses	750		
The PI-3 Kinase Pathway Is Negatively Regulated by PTEN Phosphatase	751		
		17 Cell Organization and Movement I: Microfilaments	775
16.6 Signaling Pathways Controlled by Ubiquitylation and Protein Degradation: Wnt, Hedgehog, and NF-κB	751	17.1 Microfilaments and Actin Structures	778
Wnt Signaling Triggers Release of a Transcription Factor from a Cytosolic Protein Complex	752	Actin Is Ancient, Abundant, and Highly Conserved	778
Concentration Gradients of Wnt Protein Are Essential for Many Steps in Development	753	G-Actin Monomers Assemble into Long, Helical F-Actin Polymers	779
Hedgehog Signaling Relieves Repression of Target Genes	754	F-Actin Has Structural and Functional Polarity	780
Hedgehog Signaling in Vertebrates Requires Primary Cilia	757		
Degradation of an Inhibitor Protein Activates the NF- κ B Transcription Factor	757	17.2 Dynamics of Actin Filaments	781
Polyubiquitin Chains Serve as Scaffolds Linking Receptors to Downstream Proteins in the NF- κ B Pathway	760	Actin Polymerization In Vitro Proceeds in Three Steps	781
		Actin Filaments Grow Faster at (+) Ends Than at (–) Ends	782
		Actin Filament Treadmilling Is Accelerated by Profilin and Cofilin	784
		Thymosin- β_4 Provides a Reservoir of Actin for Polymerization	785
		Capping Proteins Block Assembly and Disassembly at Actin Filament Ends	785
		17.3 Mechanisms of Actin Filament Assembly	786
		Formins Assemble Unbranched Filaments	786

The Arp2/3 Complex Nucleates Branched Filament Assembly	787
Intracellular Movements Can Be Powered by Actin Polymerization	789
Microfilaments Function in Endocytosis	790
Toxins That Perturb the Pool of Actin Monomers Are Useful for Studying Actin Dynamics	791

17.4 Organization of Actin-Based Cellular Structures 793

Cross-Linking Proteins Organize Actin Filaments into Bundles or Networks	793
Adapter Proteins Link Actin Filaments to Membranes	793

17.5 Myosins: Actin-Based Motor Proteins 796

Myosins Have Head, Neck, and Tail Domains with Distinct Functions	797
Myosins Make Up a Large Family of Mechanochemical Motor Proteins	798
Conformational Changes in the Myosin Head Couple ATP Hydrolysis to Movement	800
Myosin Heads Take Discrete Steps Along Actin Filaments	802

17.6 Myosin-Powered Movements 803

Myosin Thick Filaments and Actin Thin Filaments in Skeletal Muscle Slide Past Each Other During Contraction	803
Skeletal Muscle Is Structured by Stabilizing and Scaffolding Proteins	805
Contraction of Skeletal Muscle Is Regulated by Ca ²⁺ and Actin-Binding Proteins	805
Actin and Myosin II Form Contractile Bundles in Nonmuscle Cells	807
Myosin-Dependent Mechanisms Regulate Contraction in Smooth Muscle and Nonmuscle Cells	808
Myosin V-Bound Vesicles Are Carried Along Actin Filaments	808

17.7 Cell Migration: Mechanism, Signaling, and Chemotaxis 811

Cell Migration Coordinates Force Generation with Cell Adhesion and Membrane Recycling	811
The Small GTP-Binding Proteins Cdc42, Rac, and Rho Control Actin Organization	813
Cell Migration Involves the Coordinate Regulation of Cdc42, Rac, and Rho	815
Migrating Cells Are Steered by Chemotactic Molecules	816

18 Cell Organization and Movement II: Microtubules and Intermediate Filaments 821

18.1 Microtubule Structure and Organization 822

Microtubule Walls Are Polarized Structures Built from $\alpha\beta$ -Tubulin Dimers	822
Microtubules Are Assembled from MTOCs to Generate Diverse Configurations	824

18.2 Microtubule Dynamics 827

Individual Microtubules Exhibit Dynamic Instability	827
Localized Assembly and “Search and Capture” Help Organize Microtubules	829
Drugs Affecting Tubulin Polymerization Are Useful Experimentally and in Treatment of Diseases	829

18.3 Regulation of Microtubule Structure and Dynamics 830

Microtubules Are Stabilized by Side-Binding Proteins + TIPs Regulate the Properties and Functions of the Microtubule (+) End	831
Other End-Binding Proteins Regulate Microtubule Disassembly	832

18.4 Kinesins and Dyneins: Microtubule-Based Motor Proteins 833

Organelles in Axons Are Transported Along Microtubules in Both Directions	833
Kinesin-1 Powers Anterograde Transport of Vesicles Down Axons Toward the (+) Ends of Microtubules	835
The Kinesins Form a Large Protein Superfamily with Diverse Functions	835
Kinesin-1 Is a Highly Processive Motor	836
Dynein Motors Transport Organelles Toward the (–) Ends of Microtubules	838
Kinesins and Dyneins Cooperate in the Transport of Organelles Throughout the Cell	841
Tubulin Modifications Distinguish Different Classes of Microtubules and Their Accessibility to Motors	842

18.5 Cilia and Flagella: Microtubule-Based Surface Structures 844

Eukaryotic Cilia and Flagella Contain Long Doublet Microtubules Bridged by Dynein Motors	844
--	-----

Ciliary and Flagellar Beating Are Produced by Controlled Sliding of Outer Doublet Microtubules	844	Advancement of Neural Growth Cones Is Coordinated by Microfilaments and Microtubules	868
Intraflagellar Transport Moves Material Up and Down Cilia and Flagella	845		
Primary Cilia Are Sensory Organelles on Interphase Cells	847		
Defects in Primary Cilia Underlie Many Diseases	848		
18.6 Mitosis	849	19 The Eukaryotic Cell Cycle	873
Centrosomes Duplicate Early in the Cell Cycle in Preparation for Mitosis	849	19.1 Overview of the Cell Cycle and Its Control	875
Mitosis Can Be Divided into Six Stages	850	The Cell Cycle Is an Ordered Series of Events Leading to Cell Replication	875
The Mitotic Spindle Contains Three Classes of Microtubules	851	Cyclin-Dependent Kinases Control the Eukaryotic Cell Cycle	876
Microtubule Dynamics Increase Dramatically in Mitosis	852	Several Key Principles Govern the Cell Cycle	876
Mitotic Asters Are Pushed Apart by Kinesin-5 and Oriented by Dynein	853		
Chromosomes Are Captured and Oriented During Prometaphase	853	19.2 Model Organisms and Methods of Studying the Cell Cycle	877
Duplicated Chromosomes Are Aligned by Motors and Microtubule Dynamics	854	Budding and Fission Yeasts Are Powerful Systems for Genetic Analysis of the Cell Cycle	877
The Chromosomal Passenger Complex Regulates Microtubule Attachment at Kinetochores	855	Frog Oocytes and Early Embryos Facilitate Biochemical Characterization of the Cell Cycle Machinery	878
Anaphase A Moves Chromosomes to Poles by Microtubule Shortening	857	Fruit Flies Reveal the Interplay Between Development and the Cell Cycle	879
Anaphase B Separates Poles by the Combined Action of Kinesins and Dynein	858	The Study of Tissue Culture Cells Uncovers Cell Cycle Regulation in Mammals	880
Additional Mechanisms Contribute to Spindle Formation	858	Researchers Use Multiple Tools to Study the Cell Cycle	881
Cytokinesis Splits the Duplicated Cell in Two	859	19.3 Regulation of CDK Activity	882
Plant Cells Reorganize Their Microtubules and Build a New Cell Wall in Mitosis	860	Cyclin-Dependent Kinases Are Small Protein Kinases That Require a Regulatory Cyclin Subunit for Their Activity	883
		Cyclins Determine the Activity of CDKs	884
18.7 Intermediate Filaments	861	Cyclin Levels Are Primarily Regulated by Protein Degradation	885
Intermediate Filaments Are Assembled from Subunit Dimers	861	CDKs Are Regulated by Activating and Inhibitory Phosphorylation	886
Intermediate Filaments Are Dynamic	861	CDK Inhibitors Control Cyclin-CDK Activity	886
Cytoplasmic Intermediate Filament Proteins Are Expressed in a Tissue-Specific Manner	862	Genetically Engineered CDKs Led to the Discovery of CDK Functions	887
Lamins Line the Inner Nuclear Envelope To Provide Organization and Rigidity to the Nucleus	865		
Lamins Are Reversibly Disassembled by Phosphorylation During Mitosis	866	19.4 Commitment to the Cell Cycle and DNA Replication	887
		Cells Are Irreversibly Committed to Division at a Cell Cycle Point Called START or the Restriction Point	888
18.8 Coordination and Cooperation Between Cytoskeletal Elements	867	The E2F Transcription Factor and Its Regulator Rb Control the G ₁ -S Phase Transition in Metazoans	889
Intermediate Filament-Associated Proteins Contribute to Cellular Organization	867	Extracellular Signals Govern Cell Cycle Entry	889
Microfilaments and Microtubules Cooperate to Transport Melanosomes	867	Degradation of an S Phase CDK Inhibitor Triggers DNA Replication	890
Cdc42 Coordinates Microtubules and Microfilaments During Cell Migration	867		

Replication at Each Origin Is Initiated Once and Only Once During the Cell Cycle	892
Duplicated DNA Strands Become Linked During Replication	893

19.5 Entry into Mitosis 895

Precipitous Activation of Mitotic CDKs Initiates Mitosis	896
Mitotic CDKs Promote Nuclear Envelope Breakdown	897
Mitotic CDKs Promote Mitotic Spindle Formation	897
Chromosome Condensation Facilitates Chromosome Segregation	899

19.6 Completion of Mitosis: Chromosome Segregation and Exit from Mitosis 901

Separase-Mediated Cleavage of Cohesins Initiates Chromosome Segregation	901
APC/C Activates Separase Through Securin Ubiquitinylation	901
Mitotic CDK Inactivation Triggers Exit from Mitosis	902
Cytokinesis Creates Two Daughter Cells	903

19.7 Surveillance Mechanisms in Cell Cycle Regulation 904

Checkpoint Pathways Establish Dependencies and Prevent Errors in the Cell Cycle	905
The Growth Checkpoint Pathway Ensures That Cells Enter the Cell Cycle Only After Sufficient Macromolecule Biosynthesis	905
The DNA Damage Response System Halts Cell Cycle Progression When DNA Is Compromised	905
The Spindle Assembly Checkpoint Pathway Prevents Chromosome Segregation Until Chromosomes Are Accurately Attached to the Mitotic Spindle	908
The Spindle Position Checkpoint Pathway Ensures That the Nucleus Is Accurately Partitioned Between Two Daughter Cells	909

19.8 Meiosis: A Special Type of Cell Division 911

Extracellular and Intracellular Cues Regulate Germ Cell Formation	912
Several Key Features Distinguish Meiosis from Mitosis	912
Recombination and a Meiosis-Specific Cohesin Subunit Are Necessary for the Specialized Chromosome Segregation in Meiosis I	915
Co-orienting Sister Kinetochores Is Critical for Meiosis I Chromosome Segregation	917
DNA Replication Is Inhibited Between the Two Meiotic Divisions	917

Part IV Cell Growth and Differentiation

20 Integrating Cells into Tissues 921

20.1 Cell-Cell and Cell-Extracellular Matrix Adhesion: An Overview 923

Cell-Adhesion Molecules Bind to One Another and to Intracellular Proteins	923
The Extracellular Matrix Participates in Adhesion, Signaling, and Other Functions	925
The Evolution of Multifaceted Adhesion Molecules Made Possible the Evolution of Diverse Animal Tissues	928
Cell-Adhesion Molecules Mediate Mechanotransduction	929

20.2 Cell-Cell and Cell-Extracellular Junctions and Their Adhesion Molecules 931

Epithelial Cells Have Distinct Apical, Lateral, and Basal Surfaces	931
Three Types of Junctions Mediate Many Cell-Cell and Cell-ECM Interactions	932
Cadherins Mediate Cell-Cell Adhesions in Adherens Junctions and Desmosomes	933
Integrins Mediate Cell-ECM Adhesions, Including Those in Epithelial-Cell Hemidesmosomes	938
Tight Junctions Seal Off Body Cavities and Restrict Diffusion of Membrane Components	939
Gap Junctions Composed of Connexins Allow Small Molecules to Pass Directly Between the Cytosols of Adjacent Cells	942

20.3 The Extracellular Matrix I: The Basal Lamina 945

The Basal Lamina Provides a Foundation for Assembly of Cells into Tissues	945
Laminin, a Multi-adhesive Matrix Protein, Helps Cross-Link Components of the Basal Lamina	947
Sheet-Forming Type IV Collagen Is a Major Structural Component of the Basal Lamina	948
Perlecan, a Proteoglycan, Cross-Links Components of the Basal Lamina and Cell-Surface Receptors	950

20.4 The Extracellular Matrix II: Connective Tissue 951

Fibrillar Collagens Are the Major Fibrous Proteins in the ECM of Connective Tissues	951
Fibrillar Collagen Is Secreted and Assembled into Fibrils Outside the Cell	951
Type I and II Collagens Associate with Nonfibrillar Collagens to Form Diverse Structures	952

Proteoglycans and Their Constituent GAGs Play Diverse Roles in the ECM	953	Somatic Cells Can Generate iPS Cells	983
Hyaluronan Resists Compression, Facilitates Cell Migration, and Gives Cartilage Its Gel-Like Properties	956	ES and iPS Cells Can Generate Functional Differentiated Human Cells	986
Fibronectins Connect Cells and ECM, Influencing Cell Shape, Differentiation, and Movement	956	21.3 Stem Cells and Niches in Multicellular Organisms	987
Elastic Fibers Permit Many Tissues to Undergo Repeated Stretching and Recoiling	959	Adult Planaria Contain Pluripotent Stem Cells	988
Metalloproteases Remodel and Degrade the Extracellular Matrix	960	Multipotent Somatic Stem Cells Give Rise to Both Stem Cells and Differentiating Cells	988
20.5 Adhesive Interactions in Motile and Nonmotile Cells	961	Stem Cells for Different Tissues Occupy Sustaining Niches	988
Integrins Mediate Adhesion and Relay Signals Between Cells and Their Three-Dimensional Environment	961	Germ-Line Stem Cells Produce Sperm or Oocytes	990
Regulation of Integrin-Mediated Adhesion and Signaling Controls Cell Movement	962	Intestinal Stem Cells Continuously Generate All the Cells of the Intestinal Epithelium	991
Connections Between the ECM and Cytoskeleton Are Defective in Muscular Dystrophy	964	Hematopoietic Stem Cells Form All Blood Cells	994
IgCAMs Mediate Cell-Cell Adhesion in Neural and Other Tissues	965	Rare Types of Cells Constitute the Niche for Hematopoietic Stem Cells	996
Leukocyte Movement into Tissues Is Orchestrated by a Precisely Timed Sequence of Adhesive Interactions	966	Meristems Are Niches for Stem Cells in Plants	996
20.6 Plant Tissues	968	A Negative Feedback Loop Maintains the Size of the Shoot Apical Stem-Cell Population	998
The Plant Cell Wall Is a Laminate of Cellulose Fibrils in a Matrix of Glycoproteins	968	The Root Meristem Resembles the Shoot Meristem in Structure and Function	999
Loosening of the Cell Wall Permits Plant Cell Growth	969	21.4 Mechanisms of Cell Polarity and Asymmetric Cell Division	1000
Plasmodesmata Directly Connect the Cytosols of Adjacent Cells	969	The Intrinsic Polarity Program Depends on a Positive Feedback Loop Involving Cdc42	1000
Tunneling Nanotubes Resemble Plasmodesmata and Transfer Molecules and Organelles Between Animal Cells	970	Cell Polarization Before Cell Division Follows a Common Hierarchy of Steps	1002
Only a Few Adhesion Molecules Have Been Identified in Plants	971	Polarized Membrane Traffic Allows Yeast to Grow Asymmetrically During Mating	1003
		The Par Proteins Direct Cell Asymmetry in the Nematode Embryo	1003
		The Par Proteins and Other Polarity Complexes Are Involved in Epithelial-Cell Polarity	1007
		The Planar Cell Polarity Pathway Orients Cells Within an Epithelium	1008
		The Par Proteins Are Involved in Asymmetric Division of Stem Cells	1008
		21.5 Cell Death and Its Regulation	1011
		Most Programmed Cell Death Occurs Through Apoptosis	1012
		Evolutionarily Conserved Proteins Participate in the Apoptotic Pathway	1013
		Caspases Amplify the Initial Apoptotic Signal and Destroy Key Cellular Proteins	1015
		Neurotrophins Promote Survival of Neurons	1015
		Mitochondria Play a Central Role in Regulation of Apoptosis in Vertebrate Cells	1017
		The Pro-apoptotic Proteins Bax and Bak Form Pores and Holes in the Outer Mitochondrial Membrane	1018
21 Stem Cells, Cell Asymmetry, and Cell Death	975		
21.1 Early Mammalian Development	977		
Fertilization Unifies the Genome	977		
Cleavage of the Mammalian Embryo Leads to the First Differentiation Events	979		
21.2 Embryonic Stem Cells and Induced Pluripotent Stem Cells	980		
The Inner Cell Mass Is the Source of ES Cells	980		
Multiple Factors Control the Pluripotency of ES Cells	981		
Animal Cloning Shows That Differentiation Can Be Reversed	983		

Release of Cytochrome <i>c</i> and SMAC/DIABLO Proteins from Mitochondria Leads to Formation of the Apoptosome and Caspase Activation	1018
Trophic Factors Induce Inactivation of Bad, a Pro-apoptotic BH3-Only Protein	1018
Vertebrate Apoptosis Is Regulated by BH3-Only Pro-apoptotic Proteins That Are Activated by Environmental Stresses	1020
Two Types of Cell Murder Are Triggered by Tumor Necrosis Factor, Fas Ligand, and Related Death Signals	1021

22 Cells of the Nervous System 1025

22.1 Neurons and Glia: Building Blocks of the Nervous System 1026

Information Flows Through Neurons from Dendrites to Axons	1027
Information Moves Along Axons as Pulses of Ion Flow Called Action Potentials	1027
Information Flows Between Neurons via Synapses	1028
The Nervous System Uses Signaling Circuits Composed of Multiple Neurons	1028
Glial Cells Form Myelin Sheaths and Support Neurons	1029
Neural Stem Cells Form Nerve and Glial Cells in the Central Nervous System	1031

22.2 Voltage-Gated Ion Channels and the Propagation of Action Potentials 1034

The Magnitude of the Action Potential Is Close to E_{Na} and Is Caused by Na^+ Influx Through Open Na^+ Channels	1034
Sequential Opening and Closing of Voltage-Gated Na^+ and K^+ Channels Generate Action Potentials	1035
Action Potentials Are Propagated Unidirectionally Without Diminution	1037
Nerve Cells Can Conduct Many Action Potentials in the Absence of ATP	1039
All Voltage-Gated Ion Channels Have Similar Structures	1039
Voltage-Sensing S4 α Helices Move in Response to Membrane Depolarization	1039
Movement of the Channel-Inactivating Segment into the Open Pore Blocks Ion Flow	1042
Myelination Increases the Velocity of Impulse Conduction	1043
Action Potentials “Jump” from Node to Node in Myelinated Axons	1043
Two Types of Glia Produce Myelin Sheaths	1044
Light-Activated Ion Channels and Optogenetics	1046

22.3 Communication at Synapses 1048

Formation of Synapses Requires Assembly of Presynaptic and Postsynaptic Structures	1048
Neurotransmitters Are Transported into Synaptic Vesicles by H^+ -Linked Antiport Proteins	1052

Three Pools of Synaptic Vesicles Loaded with Neurotransmitter Are Present in the Presynaptic Terminal	1054
Influx of Ca^{2+} Triggers Release of Neurotransmitters	1054
A Calcium-Binding Protein Regulates Fusion of Synaptic Vesicles with the Plasma Membrane	1055
Fly Mutants Lacking Dynamin Cannot Recycle Synaptic Vesicles	1056
Signaling at Synapses Is Terminated by Degradation or Reuptake of Neurotransmitters	1057
Opening of Acetylcholine-Gated Cation Channels Leads to Muscle Contraction	1057
All Five Subunits in the Nicotinic Acetylcholine Receptor Contribute to the Ion Channel	1058
Nerve Cells Integrate Many Inputs to Make an All-or-None Decision to Generate an Action Potential	1059
Gap Junctions Allow Direct Communication Between Neurons and Between Glia	1060

22.4 Sensing the Environment: Touch, Pain, Taste, and Smell 1061

Mechanoreceptors Are Gated Cation Channels	1061
Pain Receptors Are Also Gated Cation Channels	1062
Five Primary Tastes Are Sensed by Subsets of Cells in Each Taste Bud	1064
A Plethora of Receptors Detect Odors	1066
Each Olfactory Receptor Neuron Expresses a Single Type of Odorant Receptor	1068

22.5 Forming and Storing Memories 1070

Memories Are Formed by Changing the Number or Strength of Synapses Between Neurons	1070
The Hippocampus Is Required for Memory Formation	1071
Multiple Molecular Mechanisms Contribute to Synaptic Plasticity	1072
Formation of Long-Term Memories Requires Gene Expression	1074

23 Immunology 1079

23.1 Overview of Host Defenses 1081

Pathogens Enter the Body Through Different Routes and Replicate at Different Sites	1081
Leukocytes Circulate Throughout the Body and Take Up Residence in Tissues and Lymph Nodes	1082
Mechanical and Chemical Boundaries Form a First Layer of Defense Against Pathogens	1083
Innate Immunity Provides a Second Line of Defense	1084
Inflammation Is a Complex Response to Injury That Encompasses Both Innate and Adaptive Immunity	1086
Adaptive Immunity, the Third Line of Defense, Exhibits Specificity	1088

23.2 Immunoglobulins: Structure and Function 1089

Immunoglobulins Have a Conserved Structure Consisting of Heavy and Light Chains	1089
Multiple Immunoglobulin Isotypes Exist, Each with Different Functions	1090
Each Naive B Cell Produces a Unique Immunoglobulin	1091
Immunoglobulin Domains Have a Characteristic Fold Composed of Two β Sheets Stabilized by a Disulfide Bond	1093
An Immunoglobulin's Constant Region Determines Its Functional Properties	1094

23.3 Generation of Antibody Diversity and B-Cell Development 1095

A Functional Light-Chain Gene Requires Assembly of V and J Gene Segments	1096
Rearrangement of the Heavy-Chain Locus Involves V, D, and J Gene Segments	1099
Somatic Hypermutation Allows the Generation and Selection of Antibodies with Improved Affinities	1099
B-Cell Development Requires Input from a Pre-B-Cell Receptor	1100
During an Adaptive Response, B Cells Switch from Making Membrane-Bound Ig to Making Secreted Ig	1101
B Cells Can Switch the Isotype of Immunoglobulin They Make	1102

23.4 The MHC and Antigen Presentation 1104

The MHC Determines the Ability of Two Unrelated Individuals of the Same Species to Accept or Reject Grafts	1104
The Killing Activity of Cytotoxic T Cells Is Antigen Specific and MHC Restricted	1105
T Cells with Different Functional Properties Are Guided by Two Distinct Classes of MHC Molecules	1105
MHC Molecules Bind Peptide Antigens and Interact with the T-Cell Receptor	1107
Antigen Presentation Is the Process by Which Protein Fragments Are Complexed with MHC Products and Posted to the Cell Surface	1109
The Class I MHC Pathway Presents Cytosolic Antigens	1110
The Class II MHC Pathway Presents Antigens Delivered to the Endocytic Pathway	1112

23.5 T Cells, T-Cell Receptors, and T-Cell Development 1115

The Structure of the T-Cell Receptor Resembles the F(ab) Portion of an Immunoglobulin	1115
TCR Genes Are Rearranged in a Manner Similar to Immunoglobulin Genes	1116

Many of the Variable Residues of TCRs Are Encoded in the Junctions Between V, D, and J Gene Segments	1118
Signaling via Antigen-Specific Receptors Triggers Proliferation and Differentiation of T and B Cells	1118
T Cells Capable of Recognizing MHC Molecules Develop Through a Process of Positive and Negative Selection	1120
T Cells Commit to the CD4 or CD8 Lineage in the Thymus	1121
T Cells Require Two Types of Signals for Full Activation	1122
Cytotoxic T Cells Carry the CD8 Co-receptor and Are Specialized for Killing	1122
T Cells Produce an Array of Cytokines That Provide Signals to Other Immune-System Cells	1123
Helper T Cells Are Divided into Distinct Subsets Based on Their Cytokine Production and Expression of Surface Markers	1124
Leukocytes Move in Response to Chemotactic Cues Provided by Chemokines	1124

23.6 Collaboration of Immune-System Cells in the Adaptive Response 1125

Toll-Like Receptors Perceive a Variety of Pathogen-Derived Macromolecular Patterns	1125
Engagement of Toll-Like Receptors Leads to Activation of Antigen-Presenting Cells	1127
Production of High-Affinity Antibodies Requires Collaboration Between B and T cells	1128
Vaccines Elicit Protective Immunity Against a Variety of Pathogens	1130
The Immune System Defends Against Cancer	1131

24 Cancer 1135

24.1 How Tumor Cells Differ from Normal Cells 1136

The Genetic Makeup of Most Cancer Cells Is Dramatically Altered	1137
Cellular Housekeeping Functions Are Fundamentally Altered in Cancer Cells	1137
Uncontrolled Proliferation Is a Universal Trait of Cancer	1139
Cancer Cells Escape the Confines of Tissues	1140
Tumors Are Heterogeneous Organs That Are Sculpted by Their Environment	1140
Tumor Growth Requires Formation of New Blood Vessels	1141
Invasion and Metastasis Are Late Stages of Tumorigenesis	1141

24.2 The Origins and Development of Cancer 1143

Carcinogens Induce Cancer by Damaging DNA	1143
Some Carcinogens Have Been Linked to Specific Cancers	1144

The Multi-hit Model Can Explain the Progress of Cancer	1145	Many Oncogenes Encode Constitutively Active Signal-Transducing Proteins	1160
Successive Oncogenic Mutations Can Be Traced in Colon Cancers	1146	Inappropriate Production of Nuclear Transcription Factors Can Induce Transformation	1160
Cancer Development Can Be Studied in Cultured Cells and in Animal Models	1146	Aberrations in Signaling Pathways That Control Development Are Associated with Many Cancers	1161
24.3 The Genetic Basis of Cancer	1149	Genes That Regulate Apoptosis Can Function as Proto-oncogenes or Tumor-Suppressor Genes	1163
Gain-of-Function Mutations Convert Proto-oncogenes into Oncogenes	1149	24.5 Deregulation of the Cell Cycle and Genome Maintenance Pathways in Cancer	1163
Cancer-Causing Viruses Contain Oncogenes or Activate Cellular Proto-oncogenes	1152	Mutations That Promote Unregulated Passage from G ₁ to S Phase Are Oncogenic	1164
Loss-of-Function Mutations in Tumor-Suppressor Genes Are Oncogenic	1152	Loss of p53 Abolishes the DNA Damage Checkpoint	1165
Inherited Mutations in Tumor-Suppressor Genes Increase Cancer Risk	1153	Loss of DNA-Repair Systems Can Lead to Cancer	1166
Epigenetic Changes Can Contribute to Tumorigenesis	1155	GLOSSARY	G-1
Micro-RNAs Can Promote and Inhibit Tumorigenesis	1155	INDEX	I-1
Researchers Are Identifying Drivers of Tumorigenesis	1156		
Molecular Cell Biology Is Changing How Cancer Is Diagnosed and Treated	1157		
24.4 Misregulation of Cell Growth and Death Pathways in Cancer	1159		
Oncogenic Receptors Can Promote Proliferation in the Absence of External Growth Factors	1159		